


Neonatal Resuscitation – 10.160



Neonatal Resuscitation – 10.160

NOTES & PRECAUTIONS:

- A. Neonatal age is birth to 28 days.
- B. Do not use atropine in neonatal resuscitation.
- C. If meconium is lightly stained and infant is vigorous (strong respiratory effort, good muscle tone, heart rate > 100 bpm) endotracheal suctioning should not be performed.
- D. An infant may need resuscitation if intrapartum risk factors for asphyxia are present (prolapsed cord, painful bleeding, prolonged rupture of membranes, maternal fever, multiple births, abnormal presentation, maternal hypotension or seizure)

KEY CONSIDERATIONS:

Fetal presentation, recent trauma, maternal health/risk factors, maternal medications, previous birth difficulties, APGAR

| APGAR SCORE: | 0 | 1 | 2 |
|--------------|-------------|-----------------------------|-----------------|
| Appearance | Blue/Pale | Body pink, blue extremities | Completely pink |
| Pulse | Absent | Slow (< 100 bpm) | ≥ 100 bpm |
| Grimace | No response | Grimace | Cough or sneeze |
| Activity | Limp | Some flexion | Active motion |
| Respirations | Absent | Slow, irregular | Good, crying |

Newborn Target Spo2 after birth:

| | |
|--------|--------|
| 1 min | 60-65% |
| 2 min | 65-70% |
| 3 min | 70-75% |
| 4 min | 75-80% |
| 5 min | 80-85% |
| 10 min | 85-95% |